

**Bryan T. Kelly, MD
Center for Hip Pain and Preservation
Hospital for Special Surgery
541 East 71st Street
New York, NY 10021**

Tel: 212.606.1159 Fax: 646.797.8865

**Hip Arthroscopy Rehabilitation
Capsular Shift with or without FAI –Labral Components**

General Guidelines:

- No external rotation greater than 30 degrees for 4 weeks
- No hyperextension for 4 weeks
- Normalize gait pattern with brace and crutches
- Weight-bearing 20 pounds
- Keep leg from rotating outward when in bed with boot or pillows
- Continuous Passive Motion Machine
 - 4 hours/day or 2 hours if on stationary bike for 2 bouts of 20-30 minute sessions

Frequency of Physical Therapy:

- Seen post-op Day 1
- Seen 1x/week for first month
- Seen 2x/week for second month
- Seen 2-3x/week for third month
- Seen 2-3x/week for fourth month

Precautions following Hip Arthroscopy/FAI: (Refixation/Osteochondroplasty)

- Weight-bearing will be determined by procedure
- Hip flexors tendonitis
- Trochanteric bursitis
- Synovitis
- Manage scarring around portal sites

Bryan T. Kelly, MD
Center for Hip Pain and Preservation
Hospital for Special Surgery
541 East 71st Street
New York, NY 10021

Tel: 212.606.1159 Fax: 646.797.8865

- Increase range of motion focusing on flexion
 - Limit external rotation and hyperextension

Guidelines:

- **Weeks 0-2**

- NO EXTERNAL ROTATION > 20 degrees
- CPM for 4 hours/day
- Bike for 20-30 minutes/day (can be 2x/day)
- Scar massage
- Hip PROM as tolerated with focus on flexion
 - IR as tolerated
 - ER max 20 deg (looking for firm end feel)
- Supine hip log rolling for internal rotation and modified external rotation
- Progress with ROM
 - Introduce stool rotations (AAROM hip IR/ER max to 20)
- Hip isometrics - NO FLEXION
 - Abduction, adduction, extension, ER
- Pelvic tilts
- Supine bridges
- NMES to quads with SAQ with pelvic tilt
- Quadruped rocking for hip flexion
- Gait training PWB with assistive device to normalize gait
- Modalities

- **Weeks 2-4**

Bryan T. Kelly, MD
Center for Hip Pain and Preservation
Hospital for Special Surgery
541 East 71st Street
New York, NY 10021

Tel: 212.606.1159 Fax: 646.797.8865

- Continue with previous therex
- Progress Weight-bearing (week 2)
 - Week 3-4: wean off crutches (2 → 1 → 0) when gait is normalized
- Progress with hip ROM
 - Bent knee fall outs (week 4)
 - Stool rotations for ER (week 3-4) max 30 degrees
 - Prone hip ER/IR (week 4)
- Glut/piriformis stretch
- Progress core strengthening (avoid hip flexor tendonitis)
- Progress with hip strengthening – isotonics all directions except flexion
 - Start isometric sub max pain free hip flexion(3-4 wks)
- Step downs
- Clam shells → isometric side-lying hip abduction
- Hip Hiking (week 4)
- Begin proprioception/balance training
 - Balance boards, single leg stance
- Bike / Elliptical (week 6)
- Scar massage
- Bilateral Cable column rotations (week 4)
- Treadmill side stepping from level surface holding on → inclines (week 5) with good abductor strength
- Aqua therapy in low end of water if available

Bryan T. Kelly, MD
Center for Hip Pain and Preservation
Hospital for Special Surgery
541 East 71st Street
New York, NY 10021

Tel: 212.606.1159 Fax: 646.797.8865

- **Weeks 4-8**

- Continue with previous therex
- Progress with ROM
 - Standing BAPS rotations
 - External rotation with FABER
 - Hip flexor, glute/piriformis, and It-band Stretching – manual and self
 - Hip flexor stretch on stool to increase hip extension
- Progress strengthening LE
 - Introduce hip flexion isotonic (Be aware of hip flexion tendonitis)
 - Multi-hip machine (open/closed chain) consider closed strength if good abductor strength
 - Leg press (bilateral → unilateral)
 - Isokinetics: knee flexion/extension
- Progress core strengthening (avoid hip flexor tendonitis)
 - Prone/side planks
- Progress with proprioception/balance
 - Bilateral → unilateral → foam → dynadisc
- Progress cable column rotations –unilateral → foam
- Side stepping with theraband
- Hip hiking on Stairmaster

- **Weeks 8-12**

- Progressive hip ROM

Bryan T. Kelly, MD
Center for Hip Pain and Preservation
Hospital for Special Surgery
541 East 71st Street
New York, NY 10021

Tel: 212.606.1159 Fax: 646.797.8865

- Hip Joint mobs with mobilization belt into limited ranges of joint motion
ONLY IF NECESSARY
 - Lateral and inferior with rotation when indicated
 - Prone posterior-anterior glides with rotation when indicated
- Progressive LE and core strengthening
- Endurance activities around the hip
- Dynamic balance activities
- **Weeks 12-16**
 - Progressive LE and core strengthening
 - Treadmill running program
 - Sport specific agility drills and plyometrics
- **3,6,12 months Re-Evaluate (Criteria for discharge)**
 - Hip Outcome Score
 - Pain free or at least a manageable level of discomfort
 - MMT within 10 percent of uninvolved LE
 - Biodex test of Quadriceps and Hamstrings peak torque within 15 percent of uninvolved
 - Single leg cross-over triple hop for distance:
 - Score of less than 85% are considered abnormal for male and female
 - Step down test